

Connecticut Department of Public Health Drug Overdose Monthly Report

Fatal Unintentional and Undetermined Drug Overdose Report

Key Findings of Drug Overdose Decedents, 2019 - October 2021

- Current monthly report is based on confirmed fatal drug overdose cases for 2019, 2020 and January through the 2nd week of October 2021. Statistics may change for 2021 as confirmation of cases occurs periodically. Period of analysis includes January 2019 through the 2nd week of October 2021.
- The final and total number of fatal drug overdoses for 2020 was 1378, with an increase of 14.6% from the previous year, 2019 with 1202 overdose deaths.
- There were 1151 confirmed and several pending cases as of the 2nd week of October 2021.
- New and emerging substances: Lethal combinations of xylazine, an animal tranquilizer, with fentanyl were identified in 2019 and continued in 2020. The same combination trends for 2021 also. In 2020, new emerging substances, Flualprazolam, a designer benzodiazepine, in combination with fentanyl resulted in 11 overdose deaths and Eutylone, a synthetic stimulant, resulted in 3 overdose deaths. Para-fluorofentanyl, a fentanyl analog, was present in 13 overdose deaths in 2020 and 78 as of the 2nd week of October 2021.

Comparative Overview of Unintentional and Undetermined Drug Overdose Deaths (January 2019 – September 2021)

	,,,
2020 and 2021	2019
There were 1,378 confirmed drug overdose	There were 1,202 unintentional and
deaths for 2020, with an increase of 14.3%	undetermined drug overdose deaths in 2019,
compared to 2019. As of the 2 nd week of October,	with an increase of 16.7% compared to 2018.
there were 1151 confirmed drug overdose deaths	
for 2021.	
The lethal combination of xylazine and fentanyl	For the first time in 2019, xylazine/fentanyl
resulted in 141 deaths in 2020 with a median	combinations were involved in 71 drug overdose
number of 11 and continues to be a problem in	deaths. The median number for xylazine-involved
2021 also. Between January and the 2 nd week of	deaths was 7.
October 2021, there were 221 deaths involving a	
fentanyl+xylazine combination.	
The average percentage of fentanyl involved	On average, fentanyl or fentanyl analogs were
deaths was at 85% for 2020 and 84.6% as of the	involved in 82% of the overdose deaths in 2019.
2 nd week of October 2021	


Updated on 11-12-2021; Data Source: Connecticut Office of the Chief Medical Examiner (OCME), per CDC-SUDORS grant guidelines.

For substance use disorder information visit: https://www.drugfreect.org.

For information on the CT DPH Opioids and Prescription Drug Overdose Prevention Program in the Office of Injury and Violence Prevention, visit: https://www.ct.gov/dph/injuryprevention


Unintentional and undetermined drug overdose deaths, by month, Connecticut, 2019 – September 2021*: There was a 14.3% increase in unintentional and undetermined drug overdose deaths in 2020 in Connecticut, compared to 2019. Confirmed drug overdose deaths in 2021 are as follows; January 130, February 115, March 122, April 120, May 138, June 126, July 144, August 143, September 100, and these numbers may increase as there are several pending cases awaiting toxicology confirmation for these months. The chart below

represents the monthly count of confirmed drug overdose deaths from January 2019 through




The percentage of fentanyl-involved overdose deaths continued to be a problem in 2021: Fentanyl-involved overdose deaths increased significantly in 2019 to 82%, compared to the previous years of 2015 to 2018. In 2020, the average percentage of fentanyl-involved deaths further increased to 85%. The percentage of fentanyl involved deaths as of the 2nd week of October 2021 was at 84.6% and it may change due to pending cases. The chart

below represents the percentage of fentanyl-involved deaths by month.




Xylazine, a veterinary tranquilizer, identified as an emerging novel substance in drug intoxication deaths in Connecticut since 2019 − September 2021*: To

enhance drug effects, recreational drugs are often adulterated with other pharmacological agents such as xylazine, a veterinary sedative not intended for human use. In Connecticut, in March 2019, xylazine was identified as a novel and emerging adulterant in fatal drug intoxications when combined with fentanyl; it continues to be a problem in 2021. There were 71 xylazine-involved deaths in 2019 and 141 in 2020. In 2021, there were 20 xylazine + fentanyl death-related events in January, 18 in February, 23 in March, 21 in April, 28 in May, 30 in June, 37 in July, 28 in August and 17 in September for a total of 221 deaths in the first seven months and these numbers may change because of pending cases. The below chart represents the number of xylazine-involved deaths.




Drug overdose death rates were higher in males compared to females in 2019 -

2020: Rates of unintentional and undetermined drug overdose-related death were consistently higher among males when compared to females. The bar graph below represents rates of unintentional and undetermined drug overdose death by gender (rate per 100,000 gender-specific population) for 2019 and 2020.




Drug overdose death rates were higher among the non-Hispanic White and non-Hispanic Black populations compared to the Hispanic population in Connecticut,

2019 – 2020: Drug overdose death rates were highest among non-Hispanic White and non-Hispanic Black populations compared to the Hispanic population during 2019 and 2020. However, in 2020, the drug overdose mortality rate substantially increased in the non-Hispanic Black population compared to the previous year. The graph below represents the unintentional and undetermined drug overdose mortality rate in Connecticut, by race/ethnicity for years 2019 and 2020.



Drug overdose death rates were highest in the 35–44-year-old age group in Connecticut, 2019 – 2020: Drug overdose death rates were calculated per 100,000 age specific population and they were highest among the 35-44 year age group followed by 25-34 and 45-54 year age groups. The graph below represents the unintentional and undetermined drug overdose mortality rate in Connecticut, by age, by year.



Drug overdose death rates in Connecticut, by County, 2019 – 2020: The graph

below represents the unintentional and undetermined drug overdose mortality rate in Connecticut, by county, by year. During this period, New Haven County (44.8) had the highest rate followed by New London County (41.9), Windham County (40.2), Litchfield County (38.8), Hartford County (37.3), and Middlesex County (39.4). In 2020, the lowest overdose death rate was seen in Fairfield County (22.0) followed by Tolland County (25.2). Overall, except Litchfield County, all other counties had an increase in drug overdose death rate in 2020, when compared to 2019.

